

First Priority Health HMO

Commercial HEDIS 2020 Results for Measurement Year 2019

| | | Measurement Year 2019 | Measurement Year 2018 | 2020 Quality Compass®** HMO National Average |
|----------------------------------|---|-----------------------|-----------------------|---|
| Prevention and Screening | | | | |
| ABA | Adult BMI Assessment | 95.24% | 91.86% | 86.34% |
| WCC | Weight Assessment and Counseling for Nutrition and Physical Activity for Children / Adolescent | | | |
| | BMI Percentile Total | 79.44% | 79.44% | 73.16% |
| | Counseling for Nutrition Total | 78.33% | 78.33% | 65.77% |
| | Counseling for Physical Activity Total | 78.06% | 78.06% | 63.48% |
| CIS | Childhood Immunization Status | | | |
| | Combo 2 - DTaP, Polio, MMR, HepB, Hib + VZV | 89.16% | 87.76% | 76.48% |
| IMA | Immunizations for Adolescents | | | |
| | Meningococcal (MCV4) | 95.05% | 93.75% | 83.03% |
| | Tdap / Td | 97.80% | 95.63% | 88.41% |
| | HPV | 30.77% | 28.75% | 36.26% |
| BCS | Breast Cancer Screening | 71.27% | 71.12% | 74.06% |
| CCS | Cervical Cancer Screening | 71.05% | 69.59% | 74.63% |
| COL | Colorectal Cancer Screening | 64.27% | 57.70% | 64.36% |
| CHL | Chlamydia Screening Combined Rate | 52.57% | 51.32% | 53.04% |
| Respiratory Conditions | | | | |
| CWP | Appropriate Testing for Pharyngitis Total ¹ | 69.44% | ND | 69.44% |
| SPR | Use of Spirometry Testing in the Assessment and Dx of COPD | 55.17% | 30.56% | 42.29% |
| PCE | Pharmacotherapy Management of COPD Exacerbation | | | |
| | Systemic Corticosteroid | <30 | <30 | 78.29% |
| | Bronchodilator | <30 | <30 | 83.40% |
| MMA | Medication Management for People With Asthma | | | |
| | Total Medication Compliance 50% | 68.82% | 72.88% | ND |
| | Total Medication Compliance 75% | 45.16% | 50.85% | 52.65% |
| AMR | Asthma Medication Ratio Total | 80.95% | 89.60% | 78.10% |
| Cardiovascular Conditions | | | | |
| CBP | Controlling High Blood Pressure | 75.43% | 75.43% | 64.29% |
| PBH | Persistence of Beta Blocker Treatment after a Heart Attack | <30 | <30 | 86.11% |
| SPC | Statin Therapy for Patients with Cardiovascular Disease | | | |
| | Received Statin Therapy - Total | 81.30% | 82.11% | 81.27% |
| | Statin Adherence 80% - Total | 84.00% | 85.15% | 75.92% |
| Diabetes | | | | |
| CDC | Comprehensive Diabetes Care | | | |
| | Hemoglobin A1c (HbA1c) Testing | 89.00% | 89.00% | 91.71% |
| | HbA1c Poor Control (>9.0%)* | 29.44% | 29.44% | 28.80% |
| | HbA1c Control (<8.0%) | 56.68% | 56.68% | 59.36% |
| | HbA1c Good Control (<7.0%) | 36.98% | 36.98% | 39.28% |
| | Diabetic Retinal Eye Exam | 59.39% | 59.39% | 58.90% |
| | Medical Attention for Nephropathy | 87.48% | 87.48% | 90.82% |
| | Blood Pressure Control (<140/90 mm Hg) | 75.13% | 75.13% | 66.55% |

First Priority Health HMO

Commercial HEDIS 2020 Results for Measurement Year 2019

| Diabetes (continued) | | | | |
|------------------------------------|---|--------|--------|--------|
| SPD | Statin Therapy for Patients With Diabetes | | | |
| | Received Statin Therapy | 63.61% | 69.85% | 65.69% |
| | Statin Adherence 80% | 78.21% | 76.50% | 70.04% |
| Behavioral Health | | | | |
| AMM | Antidepressant Medication Management | | | |
| | Acute Phase Treatment | 68.00% | 54.46% | 69.11% |
| | Continuation Phase Treatment | 58.00% | 44.55% | 52.10% |
| ADD | Follow-Up Care for Children Prescribed ADHD Medication | | | |
| | Initiation Phase | <30 | <30 | 41.62% |
| | Continuation/Maintenance Phase | <30 | <30 | 46.41% |
| FUH | Follow-up After Hospitalization for Mental Illness | | | |
| | 30 Days | 70.00% | 63.83% | 69.13% |
| | 7 Days | 47.50% | 44.68% | 49.63% |
| FUM | Follow-Up After Emergency Dept. Visit for Mental Illness | | | |
| | 30 Days | <30 | 67.65% | 59.86% |
| | 7 Days | <30 | 52.94% | 45.46% |
| FUA | Follow-Up After Emergency Dept. Visit for Alcohol and Other Drug Dependence | | | |
| | 30 Days Total | <30 | <30 | 18.02% |
| | 7 Days Total | <30 | <30 | 13.28% |
| Overuse/Appropriateness | | | | |
| URI | Appropriate Treatment for Upper Respiratory Infection ^(A) <i>higher rate indicates better performance)</i> | | | |
| | Total ¹ | 80.00% | ND | 77.64% |
| AAB | Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis* | | | |
| | Total ^{1*} | 41.94% | 25.64% | 43.55% |
| LBP | Use of Imaging Studies for Low Back Pain ^(A higher rate indicates better performance) | 80.66% | 81.36% | 79.35% |
| HDO | Use of Opioids at High Dosage * ¹ <i>The rate per 1,000 receiving Rx opioids</i> | 4.00% | 3.85% | 4.25% |
| UOP | Use of Opioids From Multiple Providers * <i>The rate per 1,000 receiving Rx opioids</i> | | | |
| | Multiple Prescribers* | 13.22% | 11.56% | 15.51% |
| | Multiple Pharmacies* | 2.20% | 2.72% | 4.67% |
| | Multiple Prescribers and Multiple Pharmacies* | 1.76% | 1.70% | 1.29% |
| Access/Availability of Care | | | | |
| IET | Initiation and Engagement of Alcohol and Other Drug Abuse or Dependence Treatment | | | |
| | Initiation Total | 44.79% | 43.75% | 34.62% |
| | Engagement Total | 14.58% | 19.79% | 11.46% |
| PPC | Prenatal and Postpartum Care ¹ | | | |
| | Timeliness of Prenatal Care | 85.71% | 91.07% | 86.25% |
| | Checkups After Delivery | 85.71% | 80.36% | 82.98% |
| CAHPS® | | | | |
| FSA | Flu Shots for Adults Ages 18 - 64 | 43.60% | 44.63% | 55.36% |

ND No data available in Quality Compass® or ND in 2019 due to first year measure; ¹Trending between Measurement Year to Prior Year should be considered with caution due to no data in prior year or change in specifications; *Lower rates indicate better performance; **National Averages taken from NCQA's Quality Compass® 2020; CAHPS® is a registered trademark of the Agency for Healthcare Research and Quality (AHRQ)