HOLECARE.

Physical Exam	Screening	Tob
at age:	Scieening	
11 Years		
12 Years	96160 Tobacco, Alcohol or Drug Use Assessment	Reco
13 Years		
14 Years		Base
15 Years		Toba
16 Years		
17 Years		of va
18 Years		
19 Years		High
20 Years		is no

Tobacco, Alcohol or Drug Use Assessment

Recommended Coding per Pennsylvania EPSDT Periodicity Schedule: 96160

Based on the Bright Futures Periodicity Schedule it is recommended that Adolescents be assessed for use of Tobacco, Alcohol or Drug use beginning at the age of 11 years old. These structured screenings require the use of validated screening tools to assess for clinical indication that the child may need further clinical evaluation.

Highmark Wholecare plans do not endorse or require any specific screening tool for screening purposes. This list is not all-inclusive, and other validated screening tools may be available. Each practice is encouraged to utilize

a screening tool that is best suited for their individual situation.

Validated Screening Tool	Age for Use	Description	Administration Time Scoring information (if available)	Where to find additional information / purchase				
	Tobacco, Alcohol or Drug Use Assessment –11 through 20 years							
Car, Relax, Alone, Forget, Friends,	12 years to 21 years	The CRAFFT is an efficient and effective health screening tool designed to identify substance use,	5 to 10 minutes	http://crafft.org/				
Trouble (CRAFFT2.1+N) English / Spanish / Vietnamese / German / Haitian Creole / Japanese / Khmer / Mandarin Chinese /		substance-related riding/driving risk, and substance use disorder among youth ages 12-21. It has been implemented as part of universal screening efforts in thousands of busy medical and community health settings, as it yields information that can serve as the basis for early intervention and patient-centered counseling.		Cost: The Suggested Tax-Deductible Donation is \$1.00/card or \$20.00/25 cards, plus \$2.50 for shipping and handling.				
Portuguese / Turkish		The CRAFFT 2.1 Screening Interview is intended for use by trained clinicians as a brief interview with an adolescent patient. The CRAFFT 2.1+N contains an additional question about tobacco and nicotine use. Self-Administered version also available						

Brief Screener for Alcohol, Tobacco, and other Drugs (BSTAD)	12 to 17 years	The BSTAD ask patients about frequency of past year use and triage them into one of three levels of substance use disorder risk: no reported use, lower risk and higher risk. Can be administered by the provider or self- administered.	2 Minutes	https://www.drugabuse.gov/nidamed- medical-health- professionals/screening-tools- resources/screening-tools-for- adolescent-substance-use Cost: Free
Screening to Brief Intervention (S2BI)	12 to 17 years	The S2BI ask patients about frequency of past year use and triage them into one of three levels of substance use disorder risk: no reported use, lower risk and higher risk. Can be administered by the provider or self- administered.	2 Minutes	https://www.drugabuse.gov/nidamed- medical-health- professionals/screening-tools- resources/screening-tools-for- adolescent-substance-use Cost: Free

Health benefits or health benefit administration may be provided by or through Highmark Wholecare, coverage by Gateway Health Plan, an independent licensee of the Blue Cross Blue Shield Association ("Highmark Wholecare").