



Clinical Guideline: The Diagnosis, Evaluation, and Treatment of Attention-Deficit/Hyperactivity Disorder (ADHD) in Children and Adolescents

Line of Business: PA Medicaid

Date of QI/UM Committee Review and Adoption: April 19, 2023

Changes for 2023	
No changes for 2023	
This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care	
Clinical Indicators	Description of the indicator
1. Follow-up Care for Children Prescribed ADHD Medication – Initiation Phase (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications, ADD)	The percentage of children newly prescribed attention-deficit/hyperactivity disorder (ADHD) medication who had at least three follow-up care visits within a 10-month period, one of which was within 30 days of when the first ADHD medication was dispensed. Two rates are reported: Initiation Phase and Continuation and Management (C&M) Phase. <i>Initiation Phase:</i> The percentage of members 6–12 years of age with a prescription dispensed for ADHD medication, who had one follow-up visit with a practitioner with prescribing authority during the 30-day Initiation Phase.
2. Follow-up for Care for Children Prescribed ADHD Medication – Continuation and Maintenance Phase (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications, ADD)	The percentage of children newly prescribed attention-deficit/hyperactivity disorder (ADHD) medication who had at least three follow-up care visits within a 10-month period, one of which was within 30 days of when the first ADHD medication was dispensed. <i>Continuation and Maintenance (C&M) Phase:</i> The percentage of members 6–12 years of age as of the IPSD with a prescription dispensed for ADHD medication, who remained on the medication for at least 210 days and who, in addition to the visit in the Initiation Phase, had at least two follow-up

	visits with a practitioner within 270 days (9 months) after the Initiation Phase ended.
References	Reference Link
AAP Clinical Practice Guideline for the Diagnosis, Evaluation, and Treatment of Attention-Deficit/Hyperactivity Disorder in Children and Adolescents (2019)	Clinical Practice Guideline for the Diagnosis, Evaluation, and Treatment of Attention-Deficit/Hyperactivity Disorder in Children and Adolescents Pediatrics American Academy of Pediatrics (aap.org)
ADHD Diagnosis and Treatment Guidelines: A Historical Perspective (2019)	ADHD Diagnosis and Treatment Guidelines: A Historical Perspective
Medscape: Pediatric Attention Deficit Hyperactivity Disorder (2022)	Pediatric Attention Deficit Hyperactivity Disorder (ADHD)
Updated ADHD guideline addresses evaluation, diagnosis, treatment from ages 4-18 (2019)	Updated ADHD Guidelines addresses evaluation, diagnosis, treatment from ages 4-18
AAP Updates Guidelines on Attention Deficit Hyperactivity Disorder with Latest Research (2019)	AAP Updates Guidelines on Attention Deficit Hyperactivity Disorder with Latest Research



Clinical Guideline: Adult Preventative Guidelines (21 & Over)

Line of Business: PA Medicaid

Date of QI/UM Committee Review and Adoption: April 19th, 2023

Changes for 2023	
<p>Added Women 16–24 years as of December 31 of the measurement year to the Chlamydia Screening description.</p> <p>Added 20 years and older as of December 31 of the measurement year to Adults’ Access to Preventive/Ambulatory Health Services description</p> <p>This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care.</p>	
Clinical Indicators	Description of the indicator
<p>1. Breast Cancer Screening (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications - <i>BCS</i>)</p>	<p>The percentage of women 50-74 years of age who had a mammogram to screen for breast cancer.</p>
<p>2. Cervical Cancer Screening (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications - <i>CCS</i>)</p>	<p>The percentage of women 21-64 years of age who were screened for cervical cancer using either of the following criteria:</p> <ul style="list-style-type: none"> • Women 21-64 years of age who had a cervical cytology performed within the last 3 years. • Women 30-64 years of age who had cervical high-risk human papillomavirus (hrHPV) testing within the last 5 years. • Women 30-64 years of age who had cervical cytology/high-risk human papillomavirus (hrHPV) co-testing performed within the last 5 years.
<p>3. Chlamydia Screening (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications - <i>CHL</i>)</p>	<p>The percentage of women 16-24 years of age who were identified as sexually active and who had at least one test for chlamydia during the measurement year.</p> <p>Women 16–24 years as of December 31 of the measurement year. Two age stratifications and a total rate are reported:</p> <ul style="list-style-type: none"> • 16-20 years • 21-24 years • Total

<p>4. Adults' Access to Preventive/Ambulatory Health Services (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications – AAP)</p>	<p>The percentage of members 20 years and older as of December 31 who had an ambulatory or preventive care visit.</p> <ul style="list-style-type: none"> • Medicaid members who had an ambulatory or preventive care visit during the measurement year. <p>Three age stratifications and a total rate are reported:</p> <ul style="list-style-type: none"> • 20-44 years • 45-64 years • 65 years and older • Total <p>The total is the sum of the age stratifications</p>
<p>Reference</p>	<p>Reference Link</p>
<p>Center for Disease Control and Prevention Recommended Adult Immunization Schedule, for Ages 19 Years and Older (2022)</p>	<p>Center for Disease Control and Prevention Recommended Adult Immunization Schedule</p>
<p>Centers for Disease Control and Prevention Promoting Health for Adults (2022)</p>	<p>Centers for Disease Control and Prevention Promoting Health for Adults</p>
<p>U.S. Preventive Task Force Recommendations Adult Preventive Health Care Schedule (2022)</p>	<p>U.S. Preventive Task Force Recommendations Adult Preventive Health Care Schedule</p>
<p>U.S. Preventive Services Task Force Final Recommendations Statement Breast Cancer: Screening (2016)</p>	<p>U.S. Preventive Services Task Force Final Recommendations Statement Breast Cancer: Screening</p>
<p>U.S. Preventive Services Task Force Final Recommendations Statement Cervical Cancer Screening (2018)</p>	<p>U.S. Preventive Services Task Force Final Recommendations Statement Cervical Cancer Screening</p>
<p>U.S. Preventive Services Task Force Final Recommendations Statement Chlamydia and Gonorrhea: Screening (2021)</p>	<p>U.S. Preventive Services Task Force Final Recommendations Statement Chlamydia and Gonorrhea: Screening</p>



Clinical Guideline: The Diagnosis and Management of Asthma

Line of Business: PA Medicare Assured

Date of QI/UM Committee Review and Adoption: April 19, 2023

Changes for 2023	
<p>No changes for 2023</p> <p>This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care</p>	
Clinical Indicators	Description of the indicator
<p>1. Asthma Medication Ratio (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications, AMR)</p>	<p>The percentage of members 5-64 years of age who were identified as having persistent asthma and had a ratio of controller medications of 0.50 or greater during the measurement year.</p> <p>Report the following age stratifications as of December 31 of the measurement year:</p> <ul style="list-style-type: none"> • 5-11 years • 12-18 years • 19-50 years • 51-64 years • Total
<p>2. Asthma in Younger Adults Admission Rate (AAR-AD) (Source: Pennsylvania Performance Measure Reporting Year (MY) 2021 Technical Specification (AAR-AD))</p>	<p>The number of discharges for asthma in enrollees ages 18 years to 39 years per 100,000 member months. Age group reported:</p> <ul style="list-style-type: none"> • 18-39 years
<p>3. Asthma in Children Admission Rate (Source: Pennsylvania Performance Measure Reporting Year (MY) 2021 Technical Specification)</p>	<p>The number of discharges for asthma in enrollees ages 2 years to 17 years per 100,000 member months. Age group reported:</p> <ul style="list-style-type: none"> • 2-17 years
References	Reference Link
<p>National Heart Lung and Blood Institute (NHLBI), National Asthma Education and Prevention Program (NAEP) (2020)</p>	<p>National Heart Lung and Blood Institute (NHLBI), National Asthma Education and Prevention Program (NAEP)</p>



Clinical Guideline: Heart Failure, MI, CAD, IVD and Cholesterol Management

Line of Business: PA Medicaid

Date of QI/UM Committee Review and Adoption: April 19, 2023

Changes for 2023	
<p>Added reference: Guideline for the Evaluation and Diagnosis of Chest Pain</p> <p>This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care</p>	
Clinical Indicators	Description of the indicator
<p>1. Persistence of Beta-Blocker Treatment after a Heart Attack (Source: HEDIS[®] Measurement Year (MY) 2023, Vol. 2, Technical Specifications - PBH)</p>	<p>The percentage of members 18 years of age and older during the measurement year who were hospitalized and discharged from July 1 of the year prior to the measurement year to June 30 of the measurement year with a diagnosis of AMI and who received persistent beta-blocker treatment for six months after discharge.</p>
<p>2. Statin Therapy for Patients with Cardiovascular Disease (Source: HEDIS[®] 2020 Measurement Year (MY), 2023, Vol. 2, Technical Specifications - SPC)</p>	<p>The percentage of males 21-75 and females 40-75 years of age during the measurement year who were identified as having clinical atherosclerotic cardiovascular disease (ASCVD) and met the following criteria: The following rates are reported:</p> <ul style="list-style-type: none"> • <i>Received statin therapy:</i> Members who were dispensed at least one high-intensity or moderate-intensity statin medication during the measurement year. • <i>Statin Adherence 80%:</i> Members who remained on a high-intensity or moderate-intensity statin medication for at least 80% of the treatment period.
References	Reference Link
<p>American College of Cardiology/American Heart Association, Task Force on Clinical Practice Guidelines (2019)</p>	<p>American College of Cardiology/American Heart Association, Task Force on Clinical Practice Guidelines</p>
<p>Journal of the American College of Cardiology, Treatment of Blood Cholesterol (2018)</p>	<p>Journal of the American College of Cardiology, Treatment of Blood Cholesterol</p>

AHA Guideline on the Management of Blood Cholesterol: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines (2018)	AHA Guideline on the Management of Blood Cholesterol: Executive Summary:
Guideline for the Management of Heart Failure (2022)	Guideline for the Management of Heart Failure
Secondary Prevention and Risk Reduction for Coronary and other Atherosclerotic Vascular Disease (2011)	Secondary Prevention and Risk Reduction for Coronary and other Atherosclerotic Vascular Disease
Addressing Social Determinants of Health in the Care of Patients with Heart Failure: A Scientific Statement from the American Heart Association (2020)	Addressing Social Determinants of Health in the Care of Patients with Heart Failure
Guideline for the Evaluation and Diagnosis of Chest Pain (2021)	Guideline for the Evaluation and Diagnosis of Chest Pain



Clinical Guideline: The Management of Chronic Obstructive Pulmonary Disease

Line of Business: PA Medicaid

Date of QI/UM Committee Review and Adoption: April 19, 2023

Changes for 2023	
<p>2023 Global Initiative for Chronic Obstructive Lung Disease (GOLD) update Added Reference: AAFP COPD: Clinical Guidance and Practice Resources</p> <p>This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care</p>	
Clinical Indicators	Description of the indicator
<p>1. Use of Spirometry Testing in the Assessment and Diagnosis of COPD (Source: HEDIS[®] Measurement Year (MY) 2023 Vol. 2, Technical Specifications - SPR)</p>	<p>Percentage of members 40 years and older with a new diagnosis of COPD or newly active COPD, who have received spirometry testing to confirm the diagnosis.</p>
<p>2. Pharmacotherapy Management of COPD Exacerbation (Source: HEDIS[®] Measurement Year (MY) 2023 Vol. 2, Technical Specifications- PCE)</p>	<p>Percentage of COPD exacerbations for members 40 years and older who had an acute inpatient discharge or ED visit (<i>any claims for COPD</i>) between January 1-November 30 of the measurement year and who were dispensed appropriate medications. Two rates are reported:</p> <ul style="list-style-type: none"> • Dispensed a systemic corticosteroid (or there was evidence of an active prescription) within 14 days of the event • Dispensed a bronchodilator (or there was evidence of an active prescription) within 30 days of the event <p>Note: The eligible population for this measure is based on acute inpatient discharges and ED visits, not on members. It is possible for the denominator to include multiple events for the same individual</p>
References	Reference Link
Global Initiative for Chronic Obstructive Lung Disease – GOLD (2023)	Global Initiative for Chronic Obstructive Lung Disease
AAFP COPD: Clinical Guidance and Practice Resources (2023)	AAFP COPD: Clinical Guidance and Practice Resources



Clinical Guideline: Cystic Fibrosis

Line of Business: PA Medicaid

Date of QI/UM Committee Review and Adoption: April 19, 2023

Changes for 2023	
No changes for 2023	
This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care	
Clinical Indicators	Description of the indicator
1. Weight Assessment and Counseling for Nutrition and Physical Activity for Children and Adolescents, (HEDIS [®] Measurement Year (MY) 2023 Vol. 2, Technical Specifications - WCC)	The percentage of members 3-17 years of age who had an outpatient visit with a PCP or OB/GYN and had evidence of the following during the measurement year: <ul style="list-style-type: none"> • BMI percentile documentation* • Counseling for nutrition • Counseling for physical activity *Because BMI norms for youth vary with age and gendersex, this measure measures BMI percentile rather than an absolute BMI value.
2. Outpatient visit with pulmonologist in the past 12 months.	Number of individuals with at least one outpatient visit with a pulmonologist in the past 12 months.
3. Annual Flu Shot	Annual flu vaccine
4. Pneumococcal Vaccine	Up-to-date on pneumococcal vaccine.
References	Reference Link
Clinical Care Guidelines, Cystic Fibrosis Foundation (2023)	Clinical Care Guidelines
Chronic Medications to Maintain Lung Health, Cystic Fibrosis Foundation (2023)	Chronic Medications to Maintain Lung Health
Age Specific Care, Cystic Fibrosis Foundation (2023)	Age Specific Care



Clinical Guideline: The Management of Major Depression in Adults in Primary Care

Line of Business: PA Medicaid

Date of QI/UM Committee Review and Adoption: April 19, 2023

Changes for 2023	
No changes for 2023	
This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care	
Clinical Indicators	Description of the indicator
1. Antidepressant Medication Management (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2 Technical Specifications- AMM)	The percentage of members 18 years of age and older who were treated with antidepressant medication, had a diagnosis of major depression and who remained on an antidepressant medication treatment. Two rates are reported: <ol style="list-style-type: none"> 1. <i>Effective Acute Phase Treatment.</i> The percentage of members who remained on an antidepressant medication for at least 84 days (12 weeks). 2. <i>Effective Continuation Phase Treatment.</i> The percentage of members who remained on an antidepressant medication for at least 180 days (6 months).
References	Reference Link
American Psychiatric Association Treating Major Depressive Disorder – A Quick Reference Guide (2010)	American Psychiatric Association Treating Major Depressive Disorder – A Quick Reference Guide
Institute for Clinical Systems Improvement Health Care, Depression, Adult Depression in Primary Care (2016)	Institute for Clinical Systems Improvement Health Care, Depression, Adult Depression in Primary Care
Agency for Healthcare Research and Quality (AHRQ), Adult Depression in Primary Care (2016)	Agency for Healthcare Research and Quality (AHRQ), Adult Depression in Primary Care

American Psychological Association Psychotherapy and Pharmacotherapy for Treating Depression (2019)	American Psychological Association Psychotherapy and Pharmacotherapy for Treating Depression
---	--



Clinical Guideline: The Management of Diabetes

Line of Business: PA Medicaid

Date of QI/UM Committee Review and Adoption: April 19, 2023

Changes for 2023	
<p>American Diabetes Association (ADA) Standards of Medical Care in Diabetes was updated for 2023 Added reference: Management of Hyperglycemia in Type 2 Diabetes (2022) Added reference: American Optometric Association, Eye Care of the Patient with Diabetes Mellitus Added reference: AHA Comprehensive Management of Cardiovascular Risk Factors for Adults with Type 2 Diabetes</p> <p>This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care</p>	
Clinical Indicators	Description of the indicator
<p>1. Hemoglobin A1c Control for Patients with Diabetes (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications, <i>HBD</i>)</p>	<p>The percentage of members 18–75 years of age with diabetes (types 1 and 2) whose hemoglobin A1c (HbA1c) <i>and non-pregnant</i> was at the following levels during the measurement year:</p> <ul style="list-style-type: none"> • HbA1c control (<8.0%). • HbA1c poor control (>9.0%). <p>Note: Organizations must use the same data collection method (Administrative or Hybrid) to report these indicators.</p>
<p>2. Eye Exam for Patients with Diabetes (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications, <i>EED</i>)</p>	<p>The percentage of members 18–75 years of age with diabetes (type 1 and type 2) who had a retinal eye exam performed.</p>
<p>3. Blood Pressure Control for Patients with Diabetes (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications, <i>BPD</i>)</p>	<p>The percentage of members 18–75 years of age with diabetes (types 1 and 2) whose blood pressure (BP) was adequately controlled (<140/90 mm Hg) during the measurement year.</p>
<p>4. Statin Therapy for Patients with Diabetes (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications, <i>SPD</i>)</p>	<p>The percentage of members 40–75 years of age during the measurement year with diabetes who do not have clinical atherosclerotic cardiovascular disease (ASCVD) who met the following criteria. Two rates are reported:</p>

	<ol style="list-style-type: none"> 1. <i>Received Statin Therapy.</i> Members who were dispensed at least one statin medication of any intensity during the measurement year 2. <i>Statin Adherence 80%.</i> Members who remained on a statin medication of any intensity for at least 80% of the treatment period.
References	Reference Link
American Diabetes Association, Standards of Medical Care (2023)	American Diabetes Association, Standards of Medical Care
Management of Hyperglycemia in Type 2 Diabetes (2022)	Management of Hyperglycemia in Type 2 Diabetes
American Optometric Association, Eye Care of the Patient with Diabetes Mellitus (2019)	American Optometric Association, Eye Care of the Patient with Diabetes Mellitus
AHA Comprehensive Management of Cardiovascular Risk Factors for Adults with Type 2 Diabetes: A Scientific Statement from the American Heart Association (2022)	AHA Comprehensive Management of Cardiovascular Risk Factors for Adults with Type 2 Diabetes



Clinical Guideline: Pediatric Preventive/EPSDT/Lead Screening - Birth to 21 Years Old

Line of Business: PA Medicaid

Date of QI/UM Committee Review and Adoption: April 19, 2023

Well Child Visits in the First 30 Months of Life

Changes for 2023	
<p>Moved the EPSDT Coding Brochure from description of the indicator to reference.</p> <p>This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care</p>	
Clinical Indicators	Description of the indicator
<p>1. Well-Child Visits in the First 30 Months of Life (Source: HEDIS® Measurement Year (MY) 2023 Vol. 2, Technical Specifications – W30)</p>	<p>The percentage of members who had the following number of well-child visits with a PCP during their last 15 months. The following rates are reported:</p> <ul style="list-style-type: none"> • Well-Child Visits in the First 15 Months Children who turned 15 months old during the measurement year: <ul style="list-style-type: none"> • Six or more well-child visits. • Well-Child Visits for Age 15 Months–30 Months. Children who turned 30 months old during the measurement year: <ul style="list-style-type: none"> • Two or more well-child visits.
<p>2. EPSDT EPSDT Visits Medicaid EPSDT Reporting Medicaid.gov</p>	<p>EPSDT Description The percentage of members who complete a visit in each periodicity. Eligibility occurs when the member is enrolled for 90 days.</p> <p>An EPSDT is identified by the use of a Well Visit Value Set (99381-99382 or 99391 -99392) with an appended EP modifier and appropriate diagnosis codes.</p>
References	Reference Link
<p>Medicaid.gov Keeping America Healthy - Early and Periodic Screening, Diagnostic, and Treatment (2022)</p>	<p>Medicaid.gov Keeping America Healthy - Early and Periodic Screening, Diagnostic, and Treatment</p>

Heath Resources & Services Administration (HRSA) Maternal & Child Health – Early Periodic Screening, Diagnostic, and Treatment (2022)	Heath Resources & Services Administration (HRSA) Maternal & Child Health – Early Periodic Screening, Diagnostic, and Treatment
EPSDT Coding Brochure (2022)	EPSDT Coding Brochure

Child and Adolescent Well-Care Visits

Changes for 2023	
<p>Moved Industry Standard from and Indicator to a Reference to align with the Well Child Visits in the First 30 Days of Life guideline. Moved EPSDT coding brochure from description of indicator to a reference</p> <p>This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care</p>	
Clinical Indicators	Description of the indicator
1. Child and Adolescent Well-Care Visits (Source: HEDIS® Measurement Year 2023 Vol. 2, Technical Specifications - WCV)	<p>The percentage of members 3–21 years of age who had at least one comprehensive well-care visit with a PCP or an OB/GYN practitioner during the measurement year.</p> <p>Report three age stratifications and total rate:</p> <ul style="list-style-type: none"> ○ 3-11 years ○ 12-17 years ○ 18-21 years ○ Total
2. EPSDT EPSDT Visits Medicaid EPSDT Reporting Medicaid.gov	<p>The percentage of members who complete a visit in each periodicity. Eligibility occurs when the member is enrolled for 90 days.</p> <p>An EPSDT is identified by the use of a Well Visit Value Set (99381-99382 or 99391 -99392) with an appended EP modifier and appropriate diagnosis codes.</p>
References	Reference Link

Medicaid.gov Keeping America Healthy - Early and Periodic Screening, Diagnostic, and Treatment (2022)	Medicaid.gov Keeping America Healthy - Early and Periodic Screening, Diagnostic, and Treatment
Heath Resources & Services Administration (HRSA) Maternal & Child Health – Early Periodic Screening, Diagnostic, and Treatment (2022)	Heath Resources & Services Administration (HRSA) Maternal & Child Health – Early Periodic Screening, Diagnostic, and Treatment
Department of Human Services (DHS) Pennsylvania’s Early and Periodic Screening, Diagnosis and Treatment (EPSDT) Program Periodicity Schedule (2022)	Pennsylvania's Early and Periodic Screening, Diagnosis and Treatment Program Periodicity Schedule
EPSDT Coding Brochure (2022)	EPSDT Coding Brochure

Child and Adolescent Immunization

Changes for 2023	
<p>Moved School Vaccination Requirements for Attendance in Pennsylvania Schools website from the description of the indicator to a reference</p> <p>This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care</p>	
Clinical Indicators	Description of the indicator
1. Childhood Immunization Status (Source: HEDIS® Measurement Year 2023 Technical Specifications - CIS)	The percentage of children 2 years of age who had four diphtheria, tetanus and acellular pertussis (DTaP); three polio (IPV); one measles, mumps and rubella (MMR); three haemophilus influenza type B (HiB); three hepatitis B (HepB), one chicken pox (VZV); four pneumococcal conjugate (PCV); one hepatitis A (HepA); two or three rotavirus (RV); and two influenza (flu) vaccines by their second birthday. The measure calculates a rate for each vaccine and three combination rates.
2. Immunizations for Adolescents (Source: HEDIS® Measurement Year (MY) 2023 Technical Specifications - IMA)	The percentage of adolescents 13 years of age who had one dose of meningococcal vaccine, one tetanus, diphtheria toxoids and acellular pertussis (Tdap) vaccine, and have completed the human papillomavirus (HPV) vaccine series by

	<p>their 13th birthday. The measure calculates a rate for each vaccine and two combination rates.</p>
<p>3. Pennsylvania EPSDT Periodicity Schedule</p>	<ul style="list-style-type: none"> • EPSDT follows the CDC and ACIP Guidelines • EPSDT tracks all HEDIS Measures, and School Vaccine Requirements from age's birth through 20 years.
<p>4. Pennsylvania Department of Health School Requirements - School Vaccination Requirements</p>	<p>KINDERGARDEN (all grade school students)</p> <ul style="list-style-type: none"> • 4 doses of tetanus, diphtheria, and acellular pertussis* (1 dose on or after the 4th birthday) • 4 doses of polio (4th dose on or after 4th birthday and at least 6 months after previous dose given)** • 2 doses of measles, mumps, rubella (MMR) • 3 doses of hepatitis B • 2 doses of varicella (chickenpox) or evidence of immunity <p>*Usually given as DTP or DTaP or DT or Td ** A fourth dose is not necessary if the third dose was administered at age 4 years or older and at least 6 months after the previous dose</p> <p>FOR ATTENDANCE IN 7TH GRADE: In addition to all other required grade school immunizations:</p> <ul style="list-style-type: none"> • 1 dose of tetanus, diphtheria, acellular pertussis (Tdap) on the first day of 7th grade. • 1 dose of meningococcal conjugate vaccine (MCV) on the first day of 7th grade. <p>Note: If a child gains entrance to school in any succeeding year, the same immunizations are required on the first day.</p> <p>FOR ATTENDANCE IN 12TH GRADE: In addition to all other required grade school immunizations:</p> <ul style="list-style-type: none"> • One dose of meningococcal conjugate vaccine (MCV) on the first day of 12th

	grade. If one dose was given at 16 years of age or older, that shall count as the 12th grade dose.
References	Reference Link
CDC Recommended Child and Adolescent Immunization Schedule ages 18 Years or Younger (2023)	CDC Recommended Child and Adolescent Immunization Schedule ages 18 Years or Younger
ACIP Vaccine-Specific Recommendations (2022)	ACIP Vaccine-Specific Recommendations
Department of Human Services (DHS) Pennsylvania’s Early and Periodic Screening, Diagnosis and Treatment (EPSDT) Program Periodicity Schedule (2022)	Pennsylvania's Early and Periodic Screening, Diagnosis and Treatment Program Periodicity Schedule
School Vaccination Requirements for Attendance in Pennsylvania Schools (2023)	School Vaccination Requirements for Attendance in Pennsylvania Schools

Developmental Screening

<p>Moved School AAP Screening Technical Assistance & Resource Center website from a clinical indicator to a reference</p> <p>Moved PA EPSDT Program Periodicity Schedule website from description of the indicator to a reference</p> <p>This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care</p>	
Clinical Indicators	Description of the indicator
<p>1. Developmental Screening in the First Three Years of Life (Source: PA EQRO Measurement Year (MY) 2022, Technical Specifications - DEV)</p>	<p>The percentage of children screened for risk of developmental, behavioral, and social delays using a standardized screening tool in the 12 months preceding or on their first, second, or third birthday. Four rates, one for each age group and a combined rate are to be calculated and reported.</p> <p>Four rates, one for each age group and a combined rate are to be calculated and reported</p> <ul style="list-style-type: none"> • Children in the eligible population who turned 1 during the measurement year. • Children in the eligible population who turned 2 during the measurement year. • Children in the eligible population who turned 3 during the measurement year.

	<ul style="list-style-type: none"> • Combined Rate <p>Children should receive Structured Developmental Screenings during the following visits:</p> <ul style="list-style-type: none"> ○ 9-11 Month Visit and ○ 18 Month Visit and ○ 30 Month Visit <p>Children Should receive a Structured Autism Screening during the following visits:</p> <ul style="list-style-type: none"> ○ 18 Month Visit and ○ 24 Month Visit <p>Children who have a positive screening should be referred for further evaluation and diagnosis without delay.</p> <p>Children must also be referred to their local Early Intervention Services. This can be done either directly through the office contacts or via CONNECT.</p>
2. Pennsylvania EPSDT Periodicity Schedule	Number of children who receive the Developmental Screening at ages listed above prior to the member turning age 2 years 9 months
References	Reference Link
PA EPSDT Program Periodicity Schedule (2022)	PA EPSDT Program Periodicity Schedule
PA Department of Education Early Learning Early Intervention (2023)	PA Department of Education Early Learning Early Intervention
AAP Screening Technical Assistance & Resource Center	AAP Screening Technical Assistance & Resource Center

Lead Testing

Changes for 2023
Moved EPSDT lead testing guidelines, environmental lead investigation, immigrant and refugee health from description of the indicator to references
Moved Medicaid lead screening and AAP blood levels among resettled refugee children in select US states 2010-2014 from description of the indicator to references

This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care	
Clinical Indicators	Description of the indicator
1. HEDIS Measure Name Lead Screening in Children (Source: HEDIS® Measurement Year 2023 Technical Specifications – LSC)	<p>The percentage of children 2 years of age who had one or more capillary or venous lead blood test for lead poisoning by their second birthday.</p> <p>All children enrolled in Medicaid are required to have 2 lead tests.</p> <ul style="list-style-type: none"> ○ Test 1 at 9-11 Months ○ Test 2 at 24 Months ○ Children who enroll in Medicaid after 24 months of age and not previously having had a lead test should receive a catch up lead test. ○ All refugee infants and children ages 0-16 should be tested for Lead. <p>For children who are refugees children should be re-tested 3 to 6 months post resettlement regardless of initial BLL result, and all children aged 6 months to 6 years should be provided with a daily pediatric multivitamin with iron.</p>
References	Reference Link
Lead Testing Guidelines Form (2022)	Lead Testing Guidelines Form
Environmental Lead Investigation Form (2022)	Environmental Lead Investigation Form
Immigrant and Refugee Health (2022)	Immigrant and Refugee Health
Medicaid.gov Lead Screening (2023)	Medicaid.gov Lead Screening
American Academy of Pediatrics, AAP Blood Lead Levels Among Resettled Refugee Children in Select US States 2010-2014 (2019)	AAP Blood Lead Levels Among Resettled Refugee Children in Select US States 2010-2014

Handling of Elevated Blood Lead Levels

Changes for 2023
<p>Moved State Regulatory Requirement EPSDT for lead levels from Summary (removed) to a clinical indicator.</p> <p>Moved Environmental Lead Investigation form from description of indicator to reference.</p> <p>Moved CDC Childhood Lead Poisoning Prevention – Healthcare Providers from description of indicator to reference.</p>

Moved CDC Recommended Actions Based on Blood Lead Level from description of indicator to reference.

This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care

Clinical Indicators	Description of the indicator
1. State Regulatory Requirement EPSDT	<ul style="list-style-type: none"> • Children who have a blood lead level of $\geq 3.5\mu\text{g}/\text{dL}$ need to receive follow up per Pennsylvania Guidelines. • Children who had an initial value $\geq 3.5\mu\text{g}/\text{dL}$ through capillary screening should have results confirmed with blood drawn by venipuncture. The CDC has a recommended schedule for obtaining a confirmatory venous sample. The higher the blood lead level on the capillary screening the more urgent the need for confirmatory testing. • Children who had an initial value of $\geq 3.5\mu\text{g}/\text{dL}$ through venipuncture should follow the CDC guidelines for follow-up blood lead testing as frequency is based on initial blood lead level. • Siblings in the home should also receive lead testing, even if previous tests showed that their lead levels were within normal limits. • Children with a venous elevated blood lead levels should be referred to Early Intervention or DART for tracking services. This can be through the CONNECT Helpline at 1-800-692-7288 • Children with a venous elevated blood lead levels should receive additional developmental screenings by their PCP to ensure that the member is achieving developmental milestones on time. • Children with a venous elevated blood lead level should be referred for an Environmental Lead Investigation on the first elevated venous lead level.
References	Reference Link
Environmental Lead Investigation Form (2022)	Environmental Lead Investigation Form
CDC Childhood Lead Poisoning Prevention – Healthcare Providers	CDC Childhood Lead Poisoning Prevention – Healthcare Providers
CDC Recommended Actions Based on Blood Lead Level	CDC Recommended Actions Based on Blood Lead Level

CDC Childhood Lead Poisoning Prevention-
Scientific Publications

[CDC Childhood Lead Poisoning Prevention-
Scientific Publications](#)



Clinical Guideline: Healthy Weight Management for Children and Adolescents

Line of Business: PA Medicaid

Date of QI/UM Committee Review and Adoption: April 19, 2023

Changes for 2023	
<p>Added reference: Prevention of Pediatric Overweight and Obesity: Position of the Academy of Nutrition and Dietetics Based on an Umbrella Review of Systematic Reviews (2022)</p> <p>Added reference: Obesity in Children and Adolescents: Screening (2017)</p> <p>Added reference: American Academy of Pediatrics Clinical Practice Guidelines for the Evaluation and Treatment of Children and Adolescents with Obesity (2023)</p> <p>This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care</p>	
Clinical Indicators	Description of the indicator
<p>1. Weight Assessment and Counseling for Nutrition and Physical Activity for Children and Adolescents, (HEDIS® Measurement Year (MY) 2023 Vol. 2, Technical Specifications - WCC)</p>	<p>1. The percentage of members 3-17 years of age who had an outpatient visit with a PCP or OB/GYN and had evidence of the following during the measurement year:</p> <ul style="list-style-type: none"> • BMI percentile documentation* • Counseling for nutrition • Counseling for physical activity <p>*Because BMI norms for youth vary with age and gender, this measure measures BMI percentile rather than an absolute BMI value</p>
References	Reference Link
CDC – Childhood Overweight and Obesity (2022)	CDC – Childhood Overweight and Obesity
PA Department of Human Services (DHS) - MA Bulletin 01-20-06 Updates to Childhood Nutrition and Weight Management Services (2020)	PA Department of Human Services (DHS) - MA Bulletin 01-20-06 Updates to Childhood Nutrition and Weight Management Services
AACE/ACE Comprehensive Clinical Practice Guidelines for Medical Care of Patients with Obesity (2016)	AACE/ACE Comprehensive Clinical Practice Guidelines for Medical Care of Patients with Obesity
JCEM – Pediatric Obesity – Assessment, Treatment and Prevention: An Endocrine Society Clinical Practice Guideline (2017)	JCEM – Pediatric Obesity – Assessment, Treatment and Prevention: An Endocrine Society Clinical Practice Guideline

Prevention of Pediatric Overweight and Obesity: Position of the Academy of Nutrition and Dietetics Based on an Umbrella Review of Systematic Reviews (2022)	Prevention of Pediatric Overweight and Obesity: Position of the Academy of Nutrition and Dietetics Based on an Umbrella Review of Systematic Reviews
Obesity in Children and Adolescents: Screening (2017)	Obesity in Children and Adolescents: Screening
American Academy of Pediatrics Clinical Practice Guidelines for the Evaluation and Treatment of Children and Adolescents with Obesity (2023)	American Academy of Pediatrics Clinical Practice Guidelines for the Evaluation and Treatment of Children and Adolescents with Obesity



Clinical Guideline: Healthy Weight Management

Line of Business: PA Medicaid

Date of QI/UM Committee Review and Adoption: April 19, 2023

Changes for 2023	
<p>Added reference: Evidence Analysis Library Adult Weight Management Guideline 2020-2021 Added reference: 2020-2025 USDA Dietary Guidelines for Americans Added reference: NIH Overweight and Obesity Treatment</p> <p>This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care</p>	
Clinical Indicators	Description of the indicator
<p>1. Obesity rates for adults in Pennsylvania by ethnicity*:</p> <ul style="list-style-type: none"> • White 32.7% • Black 44.6% • Hispanic 34.1% • Multiracial, non-Hispanic 44.9% • Asian, non-Hispanic 10.6% <p>* 2022 CDC BRFSS BMI data</p>	<p>PA Statistical Data: Age group: 18 years and older</p> <ul style="list-style-type: none"> • Racial/ethnic groups are mutually exclusive. Percentages are weighted to reflect population characteristics. • An adult who has a BMI between 25 and 29.9 is considered overweight. An adult who has a BMI of 30 or higher is considered obese. • Data based on the Behavioral Risk Factor Surveillance System, an ongoing, state-based, random-digit-dialed telephone survey of non-institutionalized civilian adults aged 18 years and older. Information about the BRFSS is available at http://www.cdc.gov/brfss/index.html. • Release date represents the date figures were accessed.
<p>2. Reduce the proportion of adults with obesity</p>	<p>Healthy People 2030 Objective: Target: 36.0 percent</p> <p>Numerator Number of adults aged 20 years and over with a body mass index (BMI) equal to or greater than 30.0</p> <p>Denominator</p>

	Number of adults aged 20 years and over
References	Reference Link
Centers for Disease Control and Prevention (CDC) – Overweight and Obesity (2022)	Centers for Disease Control and Prevention (CDC) – Overweight and Obesity
Healthy People 2030 Reduce the portion of adults with obesity (2020)	Healthy People 2030 Reduce the portion of adults with obesity
American Association of Clinical Endocrinologists and American College of Endocrinology (AACE/ACE) Clinical Practice Guidelines for Comprehensive Medical Care of Patients with Obesity (2016)	American Association of Clinical Endocrinologists and American College of Endocrinology (AACE/ACE) Clinical Practice Guidelines for Comprehensive Medical Care of Patients with Obesity
Evidence Analysis Library Adult Weight Management Guideline 2020-2021 (2021)	Evidence Analysis Library Adult Weight Management Guideline 2020-2021
2020-2025 USDA Dietary Guidelines for Americans (2020)	2020-2025 USDA Dietary Guidelines for Americans
NIH Overweight and Obesity Treatment (2022)	NIH Overweight and Obesity Treatment



Clinical Guideline: Anti-retroviral Agents in HIV-1 Infected Adults and Adolescents

Line of Business: PA Medicaid

Date of QI/UM Committee Review and Adoption: April 19, 2023

Changes for 2023	
No changes for 2023	
This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care	
Clinical Indicators	Description of the indicator
1. Outpatient visit in the past 12 months	Number of HIV+ individuals with at least one outpatient visit in the past 12 months.
2. HIV Viral Load Test during the Measurement Year – Health Resources and Services Administration (HRSA)	Percentage of enrollees age 18 and older with a diagnosis of Human Immunodeficiency Virus (HIV) who had a HIV viral load test during the measurement year. (HRSA)
3. Possession ratio of HIV medication	Percentage of individuals with pharmacy claims for HIV medications in the past 12 months with an 80% medication possession ratio.
References	Reference Link
Department of Health and Human Services (DHHS) Panel, Anti-retroviral Guidelines for Adults and Adolescents, A Working Group of the Office of AIDS Research Advisory Council (OARAC) (2022)	Guidelines for the Use of Antiretroviral Agents in Adults and Adolescents with HIV
What’s New in the COVID-19 and HIV Interim Guidance (2021)	What’s New in the COVID-19 and HIV Interim Guidance
Updated HHS Perinatal Antiretroviral Treatment Guidelines (2020)	Updated HHS Perinatal Antiretroviral Treatment Guidelines
NIH Study Finds Long-Acting Injectable Drug Prevents HIV Acquisition in Cisgender Women (2020)	NIH Study Finds Long-Acting Injectable Drug Prevents HIV Acquisition in Cisgender Women
Clinical Info HIV (2023)	Clinical Info HIV



Clinical Guideline: Prevention, Detection, Evaluation, and Treatment of High Blood Pressure

Line of Business: PA Medicaid

Date of QI/UM Committee Review and Adoption: April 19, 2023

Changes for 2023	
No changes for 2023	
This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care	
Clinical Indicators	Description of the indicator
1. Controlling High Blood Pressure (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications) <i>CBP</i>	Percentage of members 18-85 years of age who had a diagnosis of hypertension (HTN) and whose BP was adequately controlled (BP was <140/90 mm Hg) during the measurement year.
2. Controlling High Blood Pressure Ages 60 years and younger.	Percentage of members 60 years of age and younger who had a diagnosis of hypertension (HTN) and whose BP was adequately controlled (BP was <150/90 mm Hg) during the measurement year.
References	Reference Link
Journal of the American College of Cardiology, Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults (2017)	Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults
American College of Cardiology/American Heart Association, Guideline on the Primary Prevention of Cardiovascular Disease: Executive Summary (2019)	ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines
Eighth Joint National Committee (JNC 8), Management of High Blood Pressure in Adults (2014)	Management of High Blood Pressure in Adults



Clinical Guideline: Prescribing Opioids for Chronic Pain

Line of Business: PA Medicaid

Date of QI/UM Committee Review and Adoption: April 19, 2023

Changes for 2023	
<p>Added Clinical Indicator: Continued Opioid Use from HEDIS 2023</p> <p>This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care</p>	
Clinical Indicators	Description of the indicator
<p>1. Use of Opioid at High Dosage (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications - <i>HDO</i>)</p>	<p>The percentage of members 18 years and older who received prescribed opioids at a high dosage (average morphine milligram equivalent dose [MME] ≥90) for ≥15 days during the measurement year.</p> <p>Note: <i>A lower rate indicates a better performance.</i></p>
<p>2. Use of Opioids from Multiple Providers (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications - <i>UOP</i>)* <i>*Adapted with financial support from CMS and with permission from the measure developer, Pharmacy Quality Alliance (PQA).</i></p>	<p>The percentage of members 18 years and older, receiving prescription opioids for ≥15 days during the measurement year, who received opioids from multiple providers. Three rates are reported.</p> <p>1. Multiple prescribers defined as the percentage of members receiving prescriptions for opioids from four or more different prescribers during the measurement year</p> <p>2. Multiple pharmacies defined as the percentage of members receiving prescriptions for opioids from four or more different pharmacies during the measurement year.</p> <p>3. Multiple prescribers and multiple pharmacies defined as percentage of members receiving prescriptions for opioids from 4 or more different prescribers and 4 or more different pharmacies during the measurement year. (i.e., the proportion of member who are numerator compliant for both the Multiple Prescribers and</p>

	<p>Multiple Pharmacies rates).</p> <p>Note: A lower rate indicates a better performance for all three rates.</p>
<p>3 Continued Opioid Use (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications - COU)*</p> <p><i>**Adapted with financial support from the Centers for Medicare & Medicaid Services (CMS) and with permission from the measure developer, Minnesota Department of Human Services.</i></p>	<p>The percentage of members 18 years of age and older who have a new episode of opioid use that puts them at risk for continued opioid use. Two rates are reported:</p> <ol style="list-style-type: none"> 1. The percentage of members with at least 15 days of prescription opioids in a 30-day period. 2. The percentage of members with at least 31 days of prescription opioids in a 62-day period. <p>Note: A lower rate indicates better performance.</p>
References	Reference Link
CDC Guideline for Prescribing Opioid for Chronic Pain (2022)	CDC Clinical Practice Guideline for Prescribing Opioids for Pain — United States, 2022 MMWR
CDC Advises Against Misapplication of the Guideline for Prescribing Opioids for Chronic Pain (2019)	CDC Advises Against Misapplication of the Guideline for Prescribing Opioids for Chronic Pain
CDC Guideline for Prescribing Opioids for Chronic Pain-Promoting Patient Care and Safety (2021)	CDC Guideline for Prescribing Opioids for Chronic Pain-Promoting Patient Care and Safety
CDC Stacks Checklist for Prescribing Opioids for Chronic Pain (2016)	CDC Stacks Checklist for Prescribing Opioids for Chronic Pain
CDC’s Efforts to Prevent Overdoses and Substance Use-Related Harms (2022)	CDC’s Efforts to Prevent Overdoses and Substance Use-Related Harms
FDA Identifies Harm Reported from Sudden Discontinuation of Opioid Pain Medicines (2019)	FDA Identifies Harm Reported from Sudden Discontinuation of Opioid Pain Medicines
NEJM: No Shortcuts to Safer Opioid Prescribing (2019)	NEJM: No Shortcuts to Safer Opioid Prescribing



Clinical Guideline: Palliative Care

Line of Business: PA Medicaid

Date of QI/UM Committee Review and Adoption: April 19, 2023

Changes for 2023	
<p>Added to medication review clinical indicator: A medication list, signed and dated during the measurement year by the appropriate practitioner type (prescribing practitioner or clinical pharmacist), meets criteria (the practitioner's signature is considered evidence that the medications were reviewed).</p> <p>This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care</p>	
Clinical Indicators	Description of the indicator
<p>1.Care for Older Adults-Medication review (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications - COA)</p>	<p>At least one medication review conducted by a prescribing practitioner or clinical pharmacist during the measurement year and the presence of a medication list in the medical record, as documented through either administrative data or medical record review.</p> <p>A medication list, signed and dated during the measurement year by the appropriate practitioner type (prescribing practitioner or clinical pharmacist), meets criteria (the practitioner's signature is considered evidence that the medications were reviewed).</p>
<p>2.Care for Older Adults-Functional Status Assessment (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications - COA)</p>	<p>At least one functional status assessment during the measurement year, as documented through either administrative data or medical record review.</p>
<p>3.Care of the Older Adult-Pain Assessment (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications - COA)</p>	<p>At least one pain assessment during the measurement year, as documented through either administrative data or medical record review.</p>
References	Reference Link
<p>National Coalition for Hospice and Palliative Care (NCHP), National Consensus Project (NCP) Clinical Practice Guidelines for Quality Palliative Care (2018)</p>	<p>National Coalition for Hospice and Palliative Care (NCHP), National Consensus Project (NCP) Clinical Practice Guidelines for Quality Palliative Care</p>



Clinical Guideline: Pediatric Preventative Dental Care

Line of Business: PA Medicaid

Date of QI/UM Committee Review and Adoption: April 19, 2023

Changes for 2023	
<p>Retired annual dental visit clinical indicator from HEDIS Added Oral Evaluation, Dental Services (OED) as a clinical indicator from HEDIS Added Topical Fluoride for Children (TFC) as a clinical indicator from HEDIS Added Periodontal-risk assessment as a clinical indicator on the Periodicity of Examination list Added Counseling for human papilloma virus/vaccine as a clinical indicator on the Periodicity of Examination list</p> <p>This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care</p>	
Clinical Indicators	Description of the indicator
<p>1. Oral Evaluation, Dental Services (Source HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications - <i>OED</i>)</p>	<p>The percentage of members under 21 years of age who received a comprehensive or periodic oral evaluation with a dental provider during the measurement year.</p> <p>Under 21 years as of December 31 of the measurement year. Report four age stratifications and a total rate:</p> <ul style="list-style-type: none"> • 0-2 years • 3-5 years

	<ul style="list-style-type: none"> • 6-14 years • 15-20 years • Total
2. Topical Fluoride for Children (Source HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications - TFC)	<p>The percentage of members 1–4 years of age who received at least two fluoride varnish applications during the measurement year. 1–4 years as of December 31 of the measurement year. Report two age stratifications and a total rate:</p> <ul style="list-style-type: none"> • 1–2 years. • 3–4 years. • Total. <p>The total is the sum of the age stratifications.</p>
References	Reference Link
Periodicity of Examination, Preventive Dental Services, Anticipatory Guidance/Counseling, and Oral Treatment for Infants, Children, and Adolescents (2022)	Periodicity of Examination, Preventive Dental Services, Anticipatory Guidance/Counseling, and Oral Treatment for Infants, Children, and Adolescents
DHS MA Bulletin #99-21-02 PA DHS Early and Periodic Screening, Diagnosis and Treatment (EPSDT) Schedule (2021)	DHS MA Bulletin #99-21-02 PA DHS Early and Periodic Screening, Diagnosis and Treatment (EPSDT) Schedule

AGE	6-12 Months	12-24 Months	2-6 Years	6- 12 Years	12 Years and Older
Clinical Indicators					
Clinical Oral Examination ¹	✓	✓	✓	✓	✓
Assess Oral growth and development ²	✓	✓	✓	✓	✓
Caries-risk assessment ³	✓	✓	✓	✓	✓

Radiographic Assessment ⁴	✓	✓	✓	✓	✓
Prophylaxis and Topical fluoride ^{3,4}	✓	✓	✓	✓	✓
Fluoride supplementation ⁵	✓	✓	✓	✓	✓
Anticipatory guidance/counseling ⁶	✓	✓	✓	✓	✓
Oral hygiene counseling ^{3,7}	Parent	Parent	Patient/Parent	Patient/Parent	Patient
Dietary counseling ^{3,8}	✓	✓	✓	✓	✓
Counseling for nonnutritive habits ⁹	✓	✓	✓	✓	✓
Injury prevention and safety counseling ¹⁰	✓	✓	✓	✓	✓
Assess speech and language development ¹¹	✓	✓	✓		
Assessment developing occlusion ¹²			✓	✓	✓
Assessment for pit and fissure sealants ¹³			✓	✓	✓
Periodontal-risk assessment ¹⁴			✓	✓	✓
Counseling for tobacco, vaping, and substance misuse				✓	✓
Counseling for human papilloma virus/vaccine				✓	✓
Counseling for intraoral/perioral piercings				✓	✓
Assessment of third molars					✓
Transition to adult dental care					✓

¹ First examination at the eruption of the first tooth and no later than 12 months. Repeat every 6 months or as indicated by child's risk status/susceptibility to disease. Includes assessment of pathology injuries.

² By clinical examination.

³ Must be repeated regularly and frequently to maximize effectiveness.

⁴ Timing, types and frequency determined by child's history, clinical findings and susceptibility to oral disease.

⁵ Consider when systemic fluoride exposure is suboptimal. Up to at least 16 years.

⁶ Appropriate discussion and counseling should be an integral part of each visit for care.

⁷ Initially, responsibility of parent, as child matures, jointly with parent; then, when indicated, only child.

⁸ Every appointment, initially to discuss appropriate feeding practices, then the role of refined carbohydrates and frequency of snacking in caries development and childhood obesity. Monitor body mass index beginning at age two

⁹ At first, discuss the need for nonnutritive sucking: digits vs. pacifiers; then the need to wean from the habit before malocclusion or deleterious effect on the dentofacial complex occurs. For school-aged children and adolescent patients, counsel regarding any existing habits such as fingernail biting, clenching, or bruxism

¹⁰ Initially pacifiers, car seats, play objects, electric cords; secondhand smoke; when learning to walk; with sports and routine playing, including the importance of mouthguards; then motor vehicles and high-speed activities.

¹¹ Observation for age-appropriate speech articulation and fluency as well as achieving receptive and expressive language milestones.

¹² Identify: transverse, vertical, and sagittal growth patterns; asymmetry; occlusal disharmonies; functional status including temporomandibular joint dysfunction; esthetic influences on self-image and emotional development.

¹³ For caries susceptible primary molars, permanent molars, premolars, and anterior teeth with deep pits and fissures; placed as soon as possible after eruption.

¹⁴ Periodontal probing should be added to the risk-assessment process after the eruption of the first permanent molars.



Clinical Guideline: Routine and High Risk Prenatal and Postpartum Care

Line of Business: PA Medicaid

Date of QI/UM Committee Review and Adoption: April 19, 2023

Changes for 2023	
<p>No changes for 2023</p> <p>This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care</p>	
Clinical Indicators	Description of the indicator
<p>1. Timeliness of Prenatal Care (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications - PPC)</p>	<p>The percentage of deliveries of live births on or between October 8 of the year prior to the measurement year and October 7 of the measurement year. For these women, the measure assesses the following facets of prenatal and postpartum care:</p> <p><u>Timeliness of Prenatal Care.</u> The percentage of deliveries that received a prenatal care visit in the first trimester, on or before the enrollment start date or within 42 days of enrollment in the organization.</p>
<p>2. Postpartum Care (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications - PPC)</p>	<p>The percentage of deliveries of live births on or between October 8 of the year prior to the measurement year and October 7 of the measurement year. For these women, the measure assesses the following facets of prenatal and postpartum care:</p> <p><u>Postpartum Care.</u> The percentage of deliveries that had a postpartum visit on or between 7 and 84 days after delivery.</p>
<p>3. Prenatal Immunization Status (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications - PRS-E)</p>	<p>The percentage of deliveries in the Measurement Period in which women had received influenza and tetanus, diphtheria toxoids and acellular pertussis (Tdap) vaccinations.</p>

<p>4.Prenatal Depression Screening and Follow-Up (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications - <i>PND-E</i>)</p>	<p>The percentage of deliveries in which members were screened for clinical depression while pregnant and, if screened positive, received follow-up care.</p> <ol style="list-style-type: none"> 1. <i>Depression Screening</i>: The percentage of deliveries in which members were screened for clinical depression during pregnancy using a standardized instrument. 2. <i>Follow-Up on Positive Screen</i>: The percentage of deliveries in which members received follow-up care within 30 days of a positive depression screen finding.
<p>5.Postpartum Depression Screening and Follow-Up (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications -<i>PDS-E</i>)</p>	<p>The percentage of deliveries in which members were screened for clinical depression during the postpartum period, and if screened positive, received follow-up care.</p> <ol style="list-style-type: none"> 1. <i>Depression Screening</i>: The percentage of deliveries in which members were screened for clinical depression using a standardized instrument during the postpartum period. 2. <i>Follow-Up on Positive Screen</i>: The percentage of deliveries in which members received follow-up care within 30 days of a positive depression screen finding.
<p>References</p>	<p>Reference Link</p>
<p>American College of Obstetricians and Gynecologists (2021)</p>	<p>American College of Obstetricians and Gynecologists</p>
<p>National Heart, Lung and Blood Institute, Managing Asthma During Pregnancy, Pharmacologic Treatment (2004)</p>	<p>National Heart, Lung and Blood Institute, Managing Asthma During Pregnancy, Pharmacologic Treatment</p>
<p>Clinical Guidance for the Integration of the Findings of the Chronic Hypertension and Pregnancy (CHAP) Study (2022)</p>	<p>Clinical Guidance for the Integration of the Findings of the Chronic Hypertension and Pregnancy (CHAP) Study</p>
<p>American College of Allergy, Pregnancy and Asthma (2023)</p>	<p>American College of Allergy, Pregnancy and Asthma</p>
<p>Centers for Disease Control and Prevention, Depression During and After Pregnancy (2022)</p>	<p>Centers for Disease Control and Prevention, Depression During and After Pregnancy</p>

--	--



Clinical Guideline: The Treatment of Schizophrenia in Children and Adolescents

Line of Business: PA Medicaid

Date of QI/UM Committee Review and Adoption: April 19, 2023

Changes for 2023	
No changes for 2023	
This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care	
Clinical Indicators	Description of the indicator
1. Metabolic Monitoring for Children and Adolescents on Antipsychotics (Source: HEDIS® Measurement Year (MY) 2023, Volume 2 Technical Specifications, <i>APM</i>)	The percentage of children and adolescents 1-17 years of age who had two or more antipsychotic prescriptions and had metabolic testing. Three rates are reported: <ul style="list-style-type: none"> • The percentage of children and adolescents on antipsychotics who received blood glucose testing. • The percentage of children and adolescents on antipsychotics who received cholesterol testing. • The percentage of children and adolescents on antipsychotics who received blood glucose and cholesterol testing.
References	Reference Link
Journal of the American Academy of Child & Adolescent Psychiatry (JAACAP) “Practice Parameters for the Assessment and Treatment of Children and Adolescents with Schizophrenia” (2013)	Journal of the American Academy of Child & Adolescent Psychiatry (JAACAP) “Practice Parameters for the Assessment and Treatment of Children and Adolescents with Schizophrenia”
Childhood Onset Schizophrenia Treatment and Management (2019)	Childhood Onset Schizophrenia Treatment and Management



Clinical Guideline: Sickle Cell Disease

Line of Business: PA Medicaid

Date of QI/UM Committee Review and Adoption: April 19, 2023

Changes for 2023	
<p>Added new reference: National Institutes of Health, National Heart, Lung, and Blood Institute (NHLBI), Sickle Cell Disease (2022)</p> <p>Removed reference: The New England Journal of Medicine (NEJM) The Management of Sickle Cell Disease (1999)</p> <p>This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care</p>	
Clinical Indicators	Description of the indicator
1.Receipt of seasonal flu shot	Percentage of enrollees diagnosed with Sickle Cell disease who received the flu shot. (total and by race/ethnicity breakdown)
2.Receipt of meningococcal vaccination	Percentage of enrollees diagnosed with Sickle Cell disease who received the meningococcal vaccination (quadrivalent meningococcal conjugate vaccine) starting age 2-10 years, then every 5 years after)
3.Outpatient visit in the past 12 months	Percentage of enrollees diagnosed with Sickle Cell disease with at least one outpatient visit in the past 12 months.
4.Retinal Exams	Percentage of enrollees diagnosed with Sickle Cell disease who received retinal eye exams.
4.ED Visits for Pain Management	Percentage of enrollees diagnosed with Sickle Cell disease who had ED Visits for Pain Management.
5.Adherence to Antibiotic Prophylaxis	Percentage of enrollees diagnosed with Sickle Cell disease who are adherent with Antibiotic Prophylaxis.
References	Reference Link
National Institutes of Health, National Heart, Lung, and Blood Institute (NHLBI) Sickle Cell Disease (2022)	Sickle Cell Disease
National Institutes of Health, National Heart, Lung and Blood Institute (NHLBI)	Evidence-Based Management of Sickle Cell Disease: Expert Panel Report

Evidence-Based Management of Sickle Cell Disease: Expert Panel Report (2014)	
National Institutes of Health, National Heart, Lung and Blood Institute (NHLBI) The Management of Sickle Cell Disease (2002)	The Management of Sickle Cell Disease
National Library of Medicine, Quality of Care Indicators for Children with Sickle Cell Disease (2011)	Quality of Care Indicators for Children with Sickle Cell Disease
American Society of Hematology (ASH) Clinical Practice Guidelines on Sickle Cell Disease (2021)	Clinical Practice Guidelines on Sickle Cell Disease
American College of Emergency Physicians, Managing Sickle Cell Disease in the ED (2021)	Managing Sickle Cell Disease in the ED



Clinical Guideline: The Treatment of Patients with Substance Use Disorders

Line of Business: PA Medicaid

Date of QI/UM Committee Review and Adoption: April 19, 2023

Changes for 2023	
<p>No changes for 2023</p> <p>This guideline does not replace the judgment or the role of the clinician in the decision-making process for individual patients, and it is only intended to serve as an educational resource for the delivery of care</p>	
Clinical Indicators	Description of the indicator
<p>1. Initiation and Engagement of Alcohol and Other Drug Abuse or Dependence (AOD) Treatment (Source: HEDIS® Measurement Year (MY) 2023 Vol. 2, Technical Specifications, <i>IET</i>)</p>	<p>The percentage of new substance use disorder (SUD) episodes that result in treatment initiation and engagement. Two rates are reported:</p> <ol style="list-style-type: none"> <i>Initiation of SUD Treatment.</i> The percentage of new SUD episodes that result in treatment initiation through an inpatient SUD admission, outpatient visit, intensive outpatient encounter, partial hospitalization, telehealth visit or medication treatment within 14 days. <i>Engagement of SUD Treatment.</i> The percentage of new SUD episodes that have evidence of treatment engagement within 34 days of initiation.
<p>2. Follow-Up After Emergency Department Visit for Substance Use (Source: HEDIS® Measurement Year (MY) 2023, Vol. 2, Technical Specifications, <i>FUA</i>)</p> <p><i>*Adapted from an NCQA measure with financial support from the Office of the Assistant Secretary for Planning and Evaluation (ASPE) under Prime Contract No. HHSP23320100019WI/HHSP23337001T, in which NCQA was a subcontractor to Mathematica. Additional financial support was</i></p>	<p>The percentage of emergency department (ED) visits for members 13 years of age and older with a principal diagnosis of substance use disorder (SUD), or any diagnosis of drug overdose, for which there was follow-up. Two rates are reported:</p> <ol style="list-style-type: none"> The percentage of ED visits for which the member received follow-up within 30 days of the ED visit (31 total days).

<i>provided by the Substance Abuse and Mental Health Services Administration (SAMHSA).</i>	2. The percentage of ED visits for which the member received follow-up within 7 days of the ED visit (8 total days).
References	Reference Link
Practice Guideline for the Treatment of Patients with Substance Use Disorders (2010)	Practice Guideline for the Treatment of Patients with Substance Use Disorders
APA Practice Guideline for The Pharmacological Treatment of Patients with Alcohol Use Disorder (2018)	APA Practice Guideline for The Pharmacological Treatment of Patients with Alcohol Use Disorder
National Institute on Drug Abuse (NIDA) Principles of Drug Addiction Treatment: A Research-Based Guide (2018)	National Institute on Drug Abuse (NIDA) Principles of Drug Addiction Treatment: A Research-Based Guide
Dartmouth-Hitchcock Knowledge Map, Unhealthy Alcohol and Drug Use – Adult Primary Care (2017)	Dartmouth-Hitchcock Knowledge Map, Unhealthy Alcohol and Drug Use – Adult Primary Care
Dartmouth-Hitchcock Unhealthy Alcohol and Drug Use (2021)	Dartmouth-Hitchcock Unhealthy Alcohol and Drug Use
American Society of Addiction Medical (ASAM) National Practice Guideline (2020)	American Society of Addiction Medical (ASAM) National Practice Guideline
American Society of Addiction Medical (ASAM) National Practice Guideline for the Treatment of Opioid Use Disorder (2019)	American Society of Addiction Medical (ASAM) National Practice Guideline for the Treatment of Opioid Use Disorder
American Society of Addiction Medical (ASAM) Clinical Practice Guideline on Alcohol Withdrawal Management (2020)	American Society of Addiction Medical (ASAM) Clinical Practice Guideline on Alcohol Withdrawal Management